

## GlobiVac® IBH

Inclusion Body Hepatitis (IBH) Hydro Pericardium Syndrome (HPS) Vaccine, Inactivated, Serotypes FAdV-4, 8 & 11



Description: GlobiVac\* IBH is an inactivated water-in-oil emulsion vaccine containing Fowl Adenovirus Type-4 (FAdV-4), Type-8 (FAdV-8), Type-11 (FAdV-11) for active immunization against Leechi disease or Inclusion Body Hepatitis - Hydro Pericardium Syndrome.

Composition: Fach dose of 0.5 mL contains: NLT 106TCID50 of inactivated FAdV-4 virus FAdV-8 virus and FAdV-11 virus

Storage: Store in dark place at +2°C to +8°C. Protect from sunlight. Do not freeze.

Package: GlobiVac® IBH 1000 doses is packed in 500 mL HDPE bottles.

Indications: For active immunization of young chicks against Inclusion Body Hepatitis - Hydro Pericardium Syndrome (IBH-HPS) and for passive immunization in parent flocks to transfer maternal antibodies to young progeny chicks.

Vaccination schedule: Consult your Veterinarian for proper vaccination program. Birds can be vaccinated from 1st week onwards

Dosage:

Breeders & Lavers

Scenario 1: (immune the progeny by MAB)

First dose: 0.50 mL per bird during Week 12-14 weeks Second dose: 0.50 mL per bird during Week 18-20 weeks

Scenario 2: (immune the progeny by MAB in high risk zone)

First dose: 0.50 mL per bird during 12-14 weeks Second dose: 0.50 mL per bird during 18-20 weeks Third dose: 0.50 mL per bird during 40-45 weeks

Scenario 3: (immune the breeder by active immunization and the progeny by MAB)

First dose: 0.25 mL per bird during 7-10 days Second dose: 0.50 mL per bird during 12-14 weeks

Third dose: 0.50 mL per bird during 18-20 weeks

In high-risk zones, the fourth dose during 40-45 weeks for better immunization in the progeny is recommended.

 Maternal antibody (MAB) protects progeny from vertical transmission and also protects them from the disease till 4-5 weeks after birth. In very high-risk zones, it is recommended to vaccinate broilers by 0.25 mL in the first 7 days.

Broilers: 0.25 mL per bird at Day 1-7 of age.

Method of administration:

Allow the vaccine to reach ambient temperature (15°C - 25°C) before use. Shake well before administration of vaccine. Use sterile 18-20G needle and inject subcutaneously in the middle part of neck or intramuscularly in thigh muscle.

Withdrawal period: None.

Contra indications: No adverse post vaccination reaction observed in healthy birds.

## Precautions & warnings:

- Do not expose the vaccine to sunlight, heat, disinfectant and detergents.
- Do not use if breakage of emulsion is observed.
- · Do not use damaged bottles.
- Do not use unsterilized vaccination equipment.
- If accidentally injected into human being, immediate medical attention is required; take care to inform the doctor that
  the vaccine is an oil emulsion.
- Opened bottles must be used immediately.
- · Vaccinate only healthy birds, which have attained recommended body weight & age group.
- Discard all used bottles, closures and unused contents by burning.

Interactions with other medicaments & other forms of interactions: Not to be administered in combination with any other vaccines unless specified.

Undesirable effects/side effects/vaccine reaction: No clinical reaction to vaccination will be observed in healthy birds. At the site of vaccination a slight swelling may be observed after few weeks of vaccination.

Antidote for over dosing: Over dose may not cause adverse effects.

List of excipients: Span-80 & Tween-80 or Polyvac 70/Muktavac 70.

Incompatibilities: When used as per instructions, no incompatibilities reported.

Shelf life in the medical product as packaged for sale: Observe on the label.

Manufactured & Marketed by:

Globion India Private Limited,

D. No. 4-3, Sy. No. 321, Biotech Park Phase III, Karakapatla, Markook Mandal, Siddipet Dist., PIN - 502 281, Telangana, INDIA. Customer care: +91-40-27990397 | E mail: customercare@globionindia.com

Artwork Code:- F/DI/IDN/00-A/10-2021